

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



(51) International Patent Classification ⁴ : A61K 47/00, G01N 33/531		A1	(11) International Publication Number: WO 89/06976 (43) International Publication Date: 10 August 1989 (10.08.89)
(21) International Application Number: PCT/GB89/00093 (22) International Filing Date: 1 February 1989 (01.02.89) (31) Priority Application Number: 8802174 (32) Priority Date: 1 February 1988 (01.02.88) (33) Priority Country: GB (71) Applicant (for all designated States except US): QUADRANT BIORESOURCES LIMITED [GB/GB]; Manor Farm, Soulbury, Leighton Buzzard, Bedfordshire (GB). (72) Inventor; and (75) Inventor/Applicant (for US only) : ROSER, Bruce, Joseph [AU/GB]; The Old Vicarage, Church Lane, Balsham, Cambridgeshire (GB). (74) Agent: ABLEWHITE, Alan, James; Marks & Clerk, 57-60 Lincoln's Inn Fields, London WC2A 3LS (GB).			(81) Designated States: AT (European patent), AU, BE (European patent), BR, CH (European patent), DE (European patent), FR (European patent), GB, GB (European patent), HU, IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent), SU, US. Published <i>With international search report.</i> <i>With amended claims.</i>
(54) Title: METHOD OF DRYING MACROMOLECULES			
(57) Abstract			
<p>A method of drying a biological macromolecular substance by drying an aqueous solution or suspension thereof, optionally in the presence of a stabilising material, is characterised in that the aqueous solution or suspension is formulated in the presence of one or more efflorescent alkali metal, ammonium or alkaline earth metal salts and then dried.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria	FR France	ML Mali
AU Australia	GA Gabon	MR Mauritania
BB Barbados	GB United Kingdom	MW Malawi
BE Belgium	HU Hungary	NL Netherlands
BG Bulgaria	IT Italy	NO Norway
BJ Benin	JP Japan	RO Romania
BR Brazil	KP Democratic People's Republic of Korea	SD Sudan
CF Central African Republic	KR Republic of Korea	SE Sweden
CG Congo	LI Liechtenstein	SN Senegal
CH Switzerland	LK Sri Lanka	SU Soviet Union
CM Cameroon	LU Luxembourg	TD Chad
DE Germany, Federal Republic of	MC Monaco	TG Togo
DK Denmark	MG Madagascar	US United States of America
FI Finland		

Method of drying macromolecules

This invention relates to an improved process for drying biological materials, e.g. proteins and other macromolecules, viruses etc. In particular the invention concerns a method of obtaining a dried product which will remain dry without being hermetically sealed or otherwise carefully stored. The invention also relates to dried products produced by such a method.

Various methods for drying biological materials have been published. Lyophilisation of some biological materials from solutions containing various sugars has been described in GB 2009198A, GB 2126588A and JP 58074696A. The sugars include trehalose, glucose, galactose. Our earlier PCT/GB86/00396 describes the use of trehalose to stabilise such materials against drying at ambient and elevated temperatures. In addition, our UK Patent Application 88 01338 discloses the use of trehalose in the stabilisation of live viruses against freezing and drying, either frozen or at ambient temperatures, and our UK Patent Application 8715238 discloses the inclusion of trehalose in food preparations. It is thus now possible to dry safely a whole range of biological materials including

immunological components, viruses, enzymes and fluorescent agents and proteinaceous or other food stuffs.

A frequent, almost inevitable, problem with this technique is that such preparations require storage under anhydrous conditions in impermeable containers with a desiccant. This is necessary because both the macromolecules e.g. antibodies, enzymes and the like and also the carbohydrate molecules e.g. trehalose, sucrose etc. are hygroscopic when dried. In addition, the ionic buffers present usually contain deliquescent salts. All these factors combine to enhance the imbibition of water from a humid environment. This can result in the desiccated macromolecules being exposed to extremely hypertonic aqueous conditions whenever packs of dried material are opened in a humid atmosphere.

This means that such preparations are vulnerable to imperfect storage conditions and their stability can be compromised by sub-optimal handling. This occurs frequently in practice because of the tradition of keeping biologically active molecules under refrigeration. This causes condensation of atmospheric water on the cold preparation whenever the refrigerator door is opened. The preparations are also very unstable in humid or tropical climates.

This invention provides a simple method which relies on molecular mechanisms to remove imbibed water from dried macromolecular preparations, especially those protected by a sugar from desiccation damage.

According to this invention there is provided a method of drying a biological macromolecular substance by drying an aqueous solution or suspension thereof, optionally in the presence of stabilising material, characterised in that the aqueous solution or suspension is formulated in the presence of one or more efflorescent alkali metal, ammonium or alkaline earth metal salts and then dried.

The method depends on replacement of the deliquescent salts, such as sodium chloride, in conventional buffers/osmotic pressure regulators with efflorescent salts such as sodium sulphate. The term "efflorescent" is used herein to refer to salts which lose water under ambient humidity, e.g. those salts having a vapour pressure of water of crystallization of at least 15mm Hg (2000 Pa). Various salts can be used depending on the nature of the material to be dried. Naturally, the salt must be compatible with the material to be dried. For food applications the approved additive calcium lactate is efflorescent and suitable. Other efflorescent salts which might find application in certain formulations include:

1. Sodium salts: acetate trihydrate, tetraborate decahydrate (borax), bromoiridite dodecahydrate, carbonate heptahydrate, metaperiodate trihydrate, metaphosphate(tri)hexahydrate, hydrogen orthophosphate dodecahydrate, sulphite heptahydrate, thiosulphate pentahydrate (hypo).
2. Non-sodium salts (where applicable): calcium lactate hydrate, magnesium salicylate tetrahydrate, magnesium sulphate heptahydrate (Epsom salts) and ammonium sulphate.

In preparations which do not contain an added buffer, e.g. foodstuffs such as egg and milk, the efflorescent salt can be added as such, without the need to replace the buffer salt, provided that care is taken to maintain appropriate molarities.

While we do not intend to be bound by theoretical considerations, it is believed that the action of the efflorescent salt is explained as follows:

Sodium sulphate crystallises from aqueous solutions as the decahydrate $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ (Glauber's salt). Because the vapour pressure of the water of crystallisation is higher than the vapour pressure of atmospheric water even in very humid environments, this

salt actively "pumps" water from the crystals into the atmosphere. Any water imbibed by the hygroscopic desiccated preparation containing sodium sulphate crystals, is taken up by this soluble salt and in turn is expelled by the efflorescence phenomenon into the atmosphere. Thus the efflorescent salt which exists in dried preparations in a microcrystalline form, homogeneously distributed throughout the dry preparation, acts as a "molecular water pump". This maintains dried preparations in a dry state under conditions of frequent exposure to humid conditions.

When dissolved in an excess of water, sodium sulphate gives a solution of neutral pH and is non reactive in solution and non toxic to cells or tissues. A 0.12M solution is roughly isotonic with mammalian serum. It substitutes in all significant respects for the physiological salt sodium chloride.

In cases where the added salt is highly efflorescent, it is possible that the molecular water pump action will actually decrease the residual water content of the "dried" preparation. For example agarose gels containing sodium sulphate instead of sodium chloride can be dried to a clear tough film which, on storage under moderately humid atmospheric conditions, actually continue to lose water, causing blooming of the

surface as minute crystals of anhydrous sodium sulphate form. This process is entirely harmless, and the gel can be completely rehydrated.

To illustrate in practice the operation of the drying mechanism we measured actual water adsorption and desorption in a model system.

Example 1. Quantitation of water adsorption by dried protein films under controlled humidity in the presence of hygroscopic and efflorescent buffers.

Method

A 10% w/v solution of salt-free bovine plasma albumin (BSA) was made up in either physiological Dulbecco's phosphate-buffered saline (PBS) or in 0.12M sodium sulphate solution. Samples of about 1ml of solution were spread in pre-weighed sterile polystyrene tissue culture grade Petri dishes (Nunclon 60 x 15mm Cat No.1-50288) and dried by being left open in a warm room (37°C) for 48hr. They were then weighed on a 5 decimal place Mettler HK60 electronic microbalance to obtain the dry weight of protein + buffer.

They were then sealed into airtight, glass humidity chambers where they were exposed for 60hr to air in which the relative humidity was controlled at 90% with a

standard sulphuric acid solution as detailed in "The Handbook of Chemistry and Physics", 67th Edition, CRC press. They were then removed and immediately reweighed to 5 decimal places to measure the water absorbed.

All samples were then stored in an open container at room temperature and ambient relative humidity for 8 days and weighed again to measure the efficiency with which the absorbed water is desorbed.

Results. (Table I)

These show that at 90% RH much is absorbed by the dried protein. Much more is absorbed in PBS buffer (30.3%) than in sodium sulphate buffer (13.4%).

Note that the ratio of BSA to buffer in this experiment (about 1:1.7 w/w) was much higher than usually present in reagent formulations (about 1:17 w/w) so that the efficiency of the water removed by the efflorescent buffer may have been sub-optimal. In addition no trehalose or other humectant was present.

TABLE 1

Buffer	Total dry weight (mg)	Total humid weight (mg)	Water gain (%)
P.B.S	222.0	289.2	30.3
Na ₂ SO ₄	314.7	356.9	13.4

Example 2

A second type of quantitative experiment was set up with the fluorescent reagent R-Phycoerythrin (RPE) at the clinically relevant concentration of 1mg/ml in 10% w/v trehalose in either PBS or 0.12M sodium sulphate solution. The experiments were done as in the previous example.

The results (table II) show that at 90% relative humidity a large amount of water was imbibed by the dried RPE in PBS buffer where more than 20% of the dry weight of RPE was absorbed. This was not the case in sulphate buffer where only 2.4% of water was imbibed. There was a dramatic difference in the appearance of the two preparations. In the PBS buffer the RPE became liquefied and syrupy in consistency while in the sulphate buffer the dried RPE remained dry and

non-tacky. Within 4 hr of removal from the humidity chamber and storage at ambient humidity in room air, the imbibed water was desorbed from the PBS preparation to a level of 3.5-3.9%. During this process a marked phase separation occurred with large crystals of buffer separated out leaving highly concentrated syrupy protein collections in the petri dish. It is this repeated liquefaction and drying with phase separation which causes damage to the desiccated protein.

TABLE 2

Buffer	Total dry weight (mg)	Total humid weight (mg)	Water gain (%)
P.B.S	211.8	257.9	21.8
Na ₂ SO ₄	245.9	251.8	2.4

Example 3

As an example of the efficiency of this buffer system in assay kits we chose a dried preparation of a readily quantitated immunoassay system with which it was possible to measure precisely any loss of activity under sub-optimal storage conditions. The reagents consisted of a conjugate of the biotin-binding protein

Streptavidin, (purified from Streptomyces avidinii), with the fluorescent phycobiliprotein R-Phycoerythrin. This conjugate was dried in flat bottomed wells of a 96-well microtitre plate in buffers of various formulations containing the disaccharide trehalose as a means of completely preventing the loss of conjugate activity on drying.

The buffer formulations used were various ratios of standard Phosphate-buffered saline and 0.1M sodium sulphate ranging from 90% of the former plus 10% of the latter to 90% sodium sulphate buffer and 10% PBS.

Plates containing these preparations were dried in a warm room at 37°C for 24 hr. Samples of these plates were then stored at 37°C or at room temperature or at 4-10°C in a standard refrigerator for 2 months.

The dried preparations were tested for their ability to stain rat lymphocytes labelled with saturating amounts of a biotin-labelled monoclonal mouse anti-rat CD4 antibody W3/25, (D.W.Mason, R.P.Arthur, M.J.Dallman, J.R.Green, G.P.Spickett & M.L.Thomas. 1983, Immunological Reviews, 74 pp.57-82). The stained preparations were produced by rehydrating the wells of the microtitre plates with 50µl distilled water and then adding a suspension of 10^6 rat lymph-node

lymphocytes to each well. The lymphocytes were suspended in complete RPMI 1640 tissue culture medium containing 10% Haemacel as a protein supplement (TCM). Staining for 30 min at 4°C was followed by two washes in TCM before analysis in a FACS 420 Fluorescence activated cell sorter (Becton Dickinson, Sunnyvale, Calif. USA) operating an argon laser at the 488 nm line at a power of 300 mW.

The intensity of staining of these lymphocytes with the R-Phycoerythrin/Streptavidin conjugate is a direct measure of the activity of the conjugate. The fluorescence intensity is recorded as the logarithm of the brightness on the X axis against the number of cells on the Y axis. Thus any decrease in activity is revealed as a shift to the left of the modal value (peak) of the brightly fluorescent cells.

Results:-

When stored for 2 months either at room temperature or at 4-10°C in a refrigerator (Fig 1a, b) there was considerable loss of activity in buffers containing smaller amounts of 0.1M sodium sulphate buffer. This loss of activity was inversely proportional to the amount of sulphate buffer present. In 90% sulphate buffer there was no loss at all. In 50% sulphate buffer the conjugate became almost completely inactive (figs 1 and 2).

In contrast, where the conjugates were stored at 37°C in a dry warm room (Fig. 1c) the buffer formulation had little effect. There was minimal loss of activity in any of the buffer formulations.

Thus the preservation of activity in the conjugates was dependent on preservation of the dry state. This could be achieved by maintaining the conjugates in a dry environment such as the warm room or by using a high concentration of sodium sulphate buffer which maintained the dried conjugate in the dry state by efflorescence even in humid conditions.

Example 4

To show the application of this technique to a commercial product, an instant two colour immunofluorescence assay was formulated in the dry state by mixing 25µl of a solution of R-Phycoerythrin conjugated to the mouse IgG1 monoclonal anti rat CD4 antibody W3/25 and 25µl of a solution of fluorescein conjugated to the mouse IgG1 anti rat CD8 antibody MRC OX-8 (both dissolved in 0.1M sodium sulphate solution) was 50µl of 10% trehalose in distilled water in the wells of a microtitre plate and drying at 37°C. This plate was kept at room temperature for 2 months and the ability of the dry conjugates to give bright 2 colour

immunofluorescence staining with complete, specific staining of the relevant cell populations was recorded. In comparison with antibodies dried without trehalose which were almost inactive, the antibodies dried with trehalose were as good in all respects, including potency as fresh non-dried antibodies (fig. 2a, b, c).

LEGENDS

Figure 1

FACS 420 histograms of red fluorescence of Streptavidin/R-Phycoerythrin (SA/RPE) conjugate staining of W3/25 labelled rat lymphocytes. The SA/RPE conjugates were dried at 37°C in the wells of flat-bottomed microtitre plates in buffers which consisted of various mixtures of 0.1M sodium sulphate and PBS. Thereafter they were stored either at room temperature, 4-10°C in a refrigerator or at 37°C in a warm room for 2 months before use for staining.

1a. Storage at room temperature: There is progressive loss of staining potency as the proportion sulphate buffer is reduced. Curves for 90%, 70% and 50% buffers are shown. With lower proportions of sulphate buffer the conjugates were completely inactive. Storage in 90% buffer retains staining potency equal to that of fresh SA/RPE conjugate.

1b Storage at 4-10°C: Again the potency of the conjugates is greatly reduced at 50% and 70% sulphate concentration while 90% sulphate buffer retains full potency.

1c Storage at 37°C: When stored in a dry atmosphere, high proportions of sulphate buffer are not necessary to preserve staining potency.

Figure 2 Simultaneous 2-colour immunofluorescence using dried conjugates

Photographs of VDU screens of FACS 420 analysis of mixed W3/25 α CD4 antibody coupled to RPE and MRC OXB α CD8 antibody coupled to FITC on rat lymphocytes. Photographs show fluorescence on X-axis against cell number on Y-axis

F1 2 = red fluorescence on RPE

F1 1 = green fluorescence of FITC

Top: fresh conjugate

Middle: conjugate dried in 0.1M sodium sulphate containing 10% trehalose and stored for 1 month at RT⁰ before staining

Bottom: conjugate dried in 0.1M sodium sulphate buffer without trehalose showing loss of staining potency.

Sodium sulphate buffer maintains trehalose-dried conjugate with full potency when stored at room temperature but does not substitute for trehalose as a preservative of conjugate function.

Example 5

Gravimetric Tests

Method:

The model system consisted of 1ml 1% bovine serum albumin plus 10% trehalose in 10mM Trishydroxymethyl-aminomethane (Tris) chloride buffer pH 7.4 to which was added the relevant concentration of the test salts. The mixture was dried at 3° for 24hr in hydrophobic plastic dishes preweighed on a top-pan balance to an accuracy of 1 milligram, then reweighed and transferred to sealed chambers in which the relative humidity was controlled at 20-22°C by saturated solutions of various salts as described in "Handbook of chemistry and physics" 67th edition p E-42 R.C. Weast editor (1986) CRC press Paton Fla. USA. At various times the plastic dishes were removed and immediately reweighed. After about 120 hr

they were returned to ambient temperature and relative humidity and reweighed at intervals thereafter until their weight was stable.

The results are plotted as % of original weight at the various weighing times.

Other efflorescent salts tested

1. Zinc sulphate heptahydrate ($\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$)

Results: Caused protein precipitation and therefore could not be used in this application.

2. Copper sulphate pentahydrate ($\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$)

Results: Caused protein precipitation and therefore could not be used in this application.

3. Disodium hydrogen orthophosphate dodecahydrate
 $\text{Na}_2\text{PO}_4 \cdot 12\text{H}_2\text{O}$.

BSA readily dissolved in 200mM and 300mM solution. Solutions were dried at about 47% RH in an incubator room at 37° and then transferred at 24hr from the 37° room to humidity chambers at 51% RH, 80% RH or 90% RH. There was uptake of water at all 3 RH. This is probably

due to the fact that the dodecahydrate crystals lose water at 37° to the dihydrate which then absorbs water to the heptahydrate form in more humid environments. This is the form that is stable at ambient temperature and humidity. This is supported by the fact that 1ml 300mMolar disodium hydrogen orthophosphate dihydrate theoretically absorbs 1.5×10^{-3} gm moles of water to form the heptahydrate. This amounts to 27mg which is precisely the increase in weight of the phosphate-dried mixture on equilibration with ambient humidity.

The mixture increased in weight by 50% at 90% RH but only by 20% at 80% RH and about 15% at 51% RH. When transferred back to room air conditions at 144hr the adsorbed water was lost to give a stable weight at about 115-120% of initial 37° weight. Fig. 3.

4. Sodium sulphate decahydrate ($\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$).

BSA readily dissolves in 200mM or 300mM sodium sulphate. When transferred from the 37° room RH to humidity chambers, there was no water uptake at 51% or at 80% RH. There was uptake of 50% of the initial weight only at 90% RH. This water was promptly lost on return to room air. Fig. 4.

5. Sodium chloride (NaCl). [control, non efflorescent]

Mixture absorbs no water on transfer from 37° to 51% RH. Absorbs 50% of initial weight at 80% RH and 100% original weight at 90% RH. Readily loses water again when returned to room air. Fig. 5.

Summary:

Data is presented as a comparison of chloride, sulphate and phosphate on each graph at various RH in Figs. 6, 7, 8.

Both disodium hydrogen orthophosphate and sodium sulphate prevent uptake of water by the hygroscopic mixture of BSA and trehalose at 80% RH and reduce the water uptake by 50% at RH of 90% when compared with the neutral (non hygroscopic, non efflorescent) salt sodium chloride. Thus the phosphate buffer is as effective limiting the uptake of water at 90% RH when the water of hydration of the heptahydrate (stable) form of the salt is taken into account.

At 20°C these salts exert an aqueous tension equivalent to a relative humidity of:-

	Aqu tension (MmHg)	RH
NaCl	4.5	35%
$\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$	16.5	95%
$\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$	16.1	93%

Thus they permit the imbibition of water at relative humidities at or above the vapour pressures of their own water of crystallisation but not below it. for this reason Na_2SO_4 is the ideal salt as it is neutral in solution, non toxic, nonreactive with biological molecules, has a normal salty taste and will protect trehalose-dried biological molecules from moisture damage at relative humidities up to between 80 and 90%.

Disodium hydrogen orthophosphate is another example of an effective salt with a potentially slightly higher protective effect. It suffers from the disadvantage that reducing the pH will lead to the formation of sodium dihydrogen orthophosphate which is slightly deliquescent. If the pH is not adjusted a solution of disodium hydrogen phosphate in water is alkaline with a pH of about 9.5. This is thus not a suitable salt for biological molecules which require a physiological pH of 7.4.

CLAIMS:

1. A method of drying a biological macromolecular substance by drying an aqueous solution or suspension thereof, optionally in the presence of a stabilising material, characterised in that the aqueous solution or suspension is formulated in the presence of one or more efflorescent alkali metal, ammonium or alkaline earth metal salts and then dried.
2. A method according to claim 1 in which the efflorescent salt replaces sodium chloride or another hygroscopic salt as an osmotic pressure regulator and/or buffer in the solution or suspension.
3. A method according to claim 1, in which the efflorescent salt is selected from the following sodium salts:- sulphate, acetate trihydrate, tetraborate decahydrate (borax), bromoiridite dodecahydrate, carbonate heptahydrate, metaperiodate trihydrate, metaphosphate(tri)hexahydrate, hydrogen orthophosphate dodecahydrate, sulphite heptahydrate, thiosulphate pentahydrate (hypo), and the following non-sodium salts:- calcium lactate hydrate, magnesium salicylate tetrahydrate, magnesium sulphate heptahydrate (Epsom salts) and ammonium sulphate.

4. A method according to claim 1, in which the efflorescent salt is sodium sulphate or disodium hydrogen phosphate.
5. A method according to claim 1, in which the solution or suspension to be dried contains a sugar as stabilising material.
6. A method according to claim 5, in which the sugar is trehalose.

AMENDED CLAIMS

[received by the International Bureau on 21 June 1989 (21.06.89)
claims 1-2 replaced by new claim 1, claim 3 renumbered claim 2, claim 3 added (1 page)]

1. A method of drying a biological macromolecular substance by drying an aqueous solution or suspension thereof, containing one or more alkali metal, ammonium or alkaline earth metal salts as buffer or osmotic pressure regulator or the like, optionally in the presence of a stabilising material, characterised in that substantially all the said salts are selected from efflorescent salts.
2. A method according to claim 1, in which the salt is selected from the following sodium salts:- sulphate, acetate trihydrate, tetraborate decahydrate (borax), bromoiridite dodecahydrate, carbonate heptahydrate, metaperiodate trihydrate, metaphosphate(tri)hexahydrate, hydrogen orthophosphate dodecahydrate, sulphite heptahydrate, thiosulphate pentahydrate (hypo), and the following non-sodium salts:- calcium lactate hydrate, magnesium salicylate tetrahydrate, magnesium sulphate heptahydrate (Epsom salts) and ammonium sulphate.
3. A method according to claim 1, in which the salt has a vapour pressure of crystallisation of at least 15mm Hg (2000 Pa).




FIG.1a.




FIG.1b.




FIG.1c.



FIG. 2a.

FL2

FL1

FRESH ANTI-CD4 AND ANTI-CD8





FIG. 2b.

FL2

FL1

DRIED WITH PRESERVATIVE





FL2

FL1

FIG. 2c.

DRIED WITHOUT PRESERVATIVE






FIG.5.






FIG.6.



INTERNATIONAL SEARCH REPORT

International Application No PCT/GB 89/00093

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) * According to International Patent Classification (IPC) or to both National Classification and IPC IPC4: A 61 K 47/00, G 01 N 33/531																	
II. FIELDS SEARCHED <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black; margin: 5px 0;">Minimum Documentation Searched †</div> <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 30%; border-bottom: 1px solid black;">Classification System ‡</th> <th style="width: 70%; border-bottom: 1px solid black;">Classification Symbols</th> </tr> <tr> <td style="padding: 5px;">IPC4</td> <td style="padding: 5px;">A 01 N; A 61 K; G 01 N</td> </tr> </table> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black; margin: 5px 0;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched §</div>			Classification System ‡	Classification Symbols	IPC4	A 01 N; A 61 K; G 01 N											
Classification System ‡	Classification Symbols																
IPC4	A 01 N; A 61 K; G 01 N																
III. DOCUMENTS CONSIDERED TO BE RELEVANT * <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 10%; border-bottom: 1px solid black;">Category ¶</th> <th style="width: 60%; border-bottom: 1px solid black;">Citation of Document, †† with Indication, where appropriate, of the relevant passages ‡‡</th> <th style="width: 30%; border-bottom: 1px solid black;">Relevant to Claim No. ‡‡</th> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">CH, A5, 622672 (BERINGWERKE AKTIENGESELLSCHAFT) 30 April 1981, see page 3, column 2, line 15 - line 56 --</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1-4</td> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">US, A, 3539450 (A. DEUTSCH) 10 November 1970, see the whole document --</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1-5</td> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">EP, A1, 0192320 (UNILEVER PLC ET AL.) 27 August 1986, see page 4, line 32; claims 1,7 --</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1,5,6</td> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">US, A, 3862302 (R. T. PRICE ET AL.) 21 January 1975, see column 5, line 45 - line 53; abstract --</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1,2,5</td> </tr> </table>			Category ¶	Citation of Document, †† with Indication, where appropriate, of the relevant passages ‡‡	Relevant to Claim No. ‡‡	X	CH, A5, 622672 (BERINGWERKE AKTIENGESELLSCHAFT) 30 April 1981, see page 3, column 2, line 15 - line 56 --	1-4	X	US, A, 3539450 (A. DEUTSCH) 10 November 1970, see the whole document --	1-5	X	EP, A1, 0192320 (UNILEVER PLC ET AL.) 27 August 1986, see page 4, line 32; claims 1,7 --	1,5,6	X	US, A, 3862302 (R. T. PRICE ET AL.) 21 January 1975, see column 5, line 45 - line 53; abstract --	1,2,5
Category ¶	Citation of Document, †† with Indication, where appropriate, of the relevant passages ‡‡	Relevant to Claim No. ‡‡															
X	CH, A5, 622672 (BERINGWERKE AKTIENGESELLSCHAFT) 30 April 1981, see page 3, column 2, line 15 - line 56 --	1-4															
X	US, A, 3539450 (A. DEUTSCH) 10 November 1970, see the whole document --	1-5															
X	EP, A1, 0192320 (UNILEVER PLC ET AL.) 27 August 1986, see page 4, line 32; claims 1,7 --	1,5,6															
X	US, A, 3862302 (R. T. PRICE ET AL.) 21 January 1975, see column 5, line 45 - line 53; abstract --	1,2,5															
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¶ Special categories of cited documents: †‡</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p> </div> </div>																	
IV. CERTIFICATION <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; border-bottom: 1px solid black; padding: 5px;"> Date of the Actual Completion of the International Search 11th April 1989 </td> <td style="width: 50%; border-bottom: 1px solid black; padding: 5px;"> Date of Mailing of this International Search Report <div style="text-align: right; font-size: 1.2em;">27 APR 1989</div> </td> </tr> <tr> <td style="border-bottom: 1px solid black; padding: 5px;"> International Searching Authority <div style="text-align: center; font-weight: bold; margin-top: 10px;">EUROPEAN PATENT OFFICE</div> </td> <td style="border-bottom: 1px solid black; padding: 5px;"> Signature of Authorized Officer <div style="text-align: right; margin-top: 10px;"> P.C.G. VAN DER PUTTEN </div> </td> </tr> </table>			Date of the Actual Completion of the International Search 11th April 1989	Date of Mailing of this International Search Report <div style="text-align: right; font-size: 1.2em;">27 APR 1989</div>	International Searching Authority <div style="text-align: center; font-weight: bold; margin-top: 10px;">EUROPEAN PATENT OFFICE</div>	Signature of Authorized Officer <div style="text-align: right; margin-top: 10px;"> P.C.G. VAN DER PUTTEN </div>											
Date of the Actual Completion of the International Search 11th April 1989	Date of Mailing of this International Search Report <div style="text-align: right; font-size: 1.2em;">27 APR 1989</div>																
International Searching Authority <div style="text-align: center; font-weight: bold; margin-top: 10px;">EUROPEAN PATENT OFFICE</div>	Signature of Authorized Officer <div style="text-align: right; margin-top: 10px;"> P.C.G. VAN DER PUTTEN </div>																

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
X	CA, A, 1153314 (AMERICAN HOME PRODUCTS CORPORATION) 6 September 1983, see claims 1,4 --	1,2
X	Patent Abstracts of Japan, Vol 11, No 15, P536, abstract of JP 61-189454, publ 1986-08-23 (FUJISAWA PHARMACEUT CO LTD) --	1,2,5
X	EP, A2, 0030199 (MERCK & CO. INC.) 10 June 1981, see page 2, line 27; claim 1 --	1,2,5
X	GB, A, 1575155 (MERCK & CO. INC.) 17 September 1980, see page 2, line 4 - line 5; claim 1 -- -----	1,2

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO. PCT/GB 89/00093**

SA 26727

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on 03/03/89
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
CH-A5- 622672	30/04/81	NL-A- 7611152	19/04/77
		FR-A-B- 2328189	13/05/77
		DE-A- 2546166	28/04/77
		BE-A- 847354	15/04/77
		LU-A- 75987	25/05/77
		AU-D- 18694/76	20/04/78
		US-A- 4157383	05/06/79
		GB-A- 1554886	31/10/79
		JP-A- 52047916	16/04/77
		AT-A- 355228	25/02/80
		CA-A- 1077394	13/05/80
		AU-A- 511176	31/07/80
		SE-A- 7611505	16/04/77
US-A- 3539450	10/11/70	NL-A- 6708570	02/01/68
		GB-A- 1163409	04/09/69
		GB-A- 1191697	13/05/70
		GB-A- 1192046	13/05/70
		US-A- 3527332	08/09/70
		US-A- 3527331	08/09/70
		US-A- 3527674	08/09/70
		US-A- 3540984	17/11/70
		US-A- 3539453	10/11/70
		BE-A- 700778	02/01/68
		CH-A- 514841	31/10/71
		DE-A- 1598321	24/02/72
		DE-A- 1598326	02/03/72
		DE-A- 1598325	11/01/73
EP-A1- 0192320	27/08/86	WO-A- 86/04095	17/07/86
US-A- 3862302	21/01/75	NL-A- 7003963	22/09/70
		DE-A- 2013498	24/09/70
		FR-A- 2037234	31/12/70
		US-A- 3655838	11/04/72
		GB-A- 1304045	24/01/73
		CH-A- 546570	15/03/74
		BE-A- 747699	21/09/70
		SE-A-C- 376087	05/05/75
CA-A- 1153314	06/09/83	US-A- 4295280	20/10/81
EP-A2- 0030199	10/06/81	US-A- 4273762	16/06/81
		JP-A- 56092822	27/07/81
		CA-A- 1146861	24/05/83

EPO FORM P007

ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.

PCT/GB 89/00093

SA 26727

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 03/03/89. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A2- 0030199	10/06/81	AT-E- 6201	84/03/15
GB-A- 1575155	17/09/80	NONE	